

# ICD-10 Preparation

2015 Weblab Series

January / Billers

# ICD-10 Roadmap

- January 25<sup>th</sup>
  - v6.26
  - Referral and EHR cleanup
- March 15<sup>th</sup>
  - v6.27

# March Release

- Version 6.27
  - ICD10 to ICD9 in Partner

# Pediatric ICD-9s and SNOMEDs We've Converted

- Asthma
- Fever
- Immunizations
- Influenza
- Pharyngitis
- Rhinitis
- Shaken baby syndrome
- UTI
- URI
- Viral Syndrome

# Otitis Media

- ICD-9 - Acute
  - 381.00 Acute *nonsuppurative* otitis media, **unspecified**
  - 381.01 Acute *serous* otitis media
  - 381.02 Acute *muroid* otitis media
  - 381.03 Acute *sanguinous* otitis media
  - 381.04 Acute *allergic serous* otitis media
  - 381.05 Acute *allergic muroid* otitis media
  - 381.06 Acute *allergic sanguinous* otitis media

# Otitis Media

- ICD-9 – Acute (cont'd)
  - 382.00 Acute suppurative otitis media *without* spontaneous rupture of eardrum
  - 382.01 Acute suppurative otitis media *with* spontaneous rupture of eardrum
  - 382.02 Acute suppurative otitis media in diseases classified elsewhere

# Otitis Media

- ICD-9 – Chronic
  - 381.10 Chronic serous otitis media, *simple* or unspecified
  - 381.19 Other chronic serous otitis media
  - 381.20 Chronic mucoid otitis media, simple or unspecified
  - 381.29 Other chronic mucoid otitis media

# Otitis Media

- ICD-9 – Chronic (cont'd)
  - 381.3 Other and *unspecified* chronic nonsuppurative otitis media
  - 382.1 Chronic *tubotympanic suppurative* otitis media
  - 382.2 Chronic *atticoantral suppurative* otitis media



# Otitis Media – ICD-10

## **Diseases of middle ear and mastoid (H65-H75)**

### **H65 Nonsuppurative otitis media**

**Includes:** nonsuppurative otitis media with myringitis

**Use additional code** for any associated perforated tympanic membrane (H72.-)

**Use additional code to identify:**

exposure to environmental tobacco smoke (Z77.22)

exposure to tobacco smoke in the perinatal period (P96.81)

history of tobacco use (Z87.891)

occupational exposure to environmental tobacco smoke (Z57.31)

tobacco dependence (F17.-)

tobacco use (Z72.0)

# Otitis Media – ICD-10

- Laterality: Left, right, bilateral, **unspecified**
- Occurrence: Recurrent, acute, chronic
- Types: Allergic (acute, subacute), serous, nonsuppurative, mucoid, sanguinous, seromucinous, tubotympanic, atticofacial
- Rupture:
  - Without
  - With: code for perforation too

# Otitis Media – ICD-10

## **Otitis media in diseases classified elsewhere**

**Code first** underlying disease, such as:  
viral disease NEC (B00-B34)

**Use additional** code for any associated perforated tympanic membrane (H72.-)

**Excludes1:** otitis media in:

influenza (J09.X9, J10.83, J11.83)

measles (B05.3)

scarlet fever (A38.0)

tuberculosis (A18.6)

# January 2015 AAP Coding NL Highlights

- Revised ICD-10 guidelines 2015
  - Clarity on Initial vs. subsequent encounters
- ICD-10 concept: Late effects and sequelae
- Advice for using new “X series” of -59 modifiers

# -59 Subset Modifiers

- Services are distinct because...:
  - XE ...it occurred during a separate encounter
  - XS ...it was performed on a separate organ/structure
  - XP ... it was performed by a different practitioner
  - XU ...it does not overlap usual components of the main service

# Suggestions? Questions?

*258* days until Thursday, 10/1/15

Next Weblab:

**Fri 2/13/15 12:00p EST**

# Resources

- Slides, reference materials, links, video recordings:
  - <http://www.pedsources.com/ICD-10-resources>
- 2015 ICD-10 Guidelines:
  - [http://www.cdc.gov/nchs/data/icd/icd10cm\\_guidelines\\_2015.pdf](http://www.cdc.gov/nchs/data/icd/icd10cm_guidelines_2015.pdf)

# Many Thanks for Attending!

[support@pcc.com](mailto:support@pcc.com)

[www.pedsource.com](http://www.pedsource.com)