

CMS ID & CQM NO.	CQM TITLE	DESCRIPTION
CMS146v1 NQF 0002	Appropriate Testing for Children with Pharyngitis	Percentage of children 2-18 years of age, who were diagnosed with pharyngitis, ordered an antibiotic and received a group A streptococcus (strep) test for the episode.
CMS155v1 NQF 0024	Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents	<p>Percentage of patients 3-17 years of age who had an outpatient visit with a Primary Care Physician (PCP) or Obstetrician/Gynecologist (OB/GYN) and who had evidence of the following during the measurement period.</p> <p>Three rates are reported.</p> <ul style="list-style-type: none"> • Percentage of patients with height, weight, and body mass index (BMI) percentile documentation. • Percentage of patients with counseling for nutrition. • Percentage of patients with counseling for physical activity.
CMS153v1 NQF 0033	Chlamydia Screening for Women	Percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement period.
CMS126v1 NQF 0036	Use of Appropriate Medications for Asthma	Percentage of patients 5-64 years of age who were identified as having persistent asthma and were appropriately prescribed medication during the measurement period.
MS117v1 NQF 0038	Childhood Immunization Status	Percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV), one measles, mumps and rubella (MMR); three H influenza type B (HiB); three hepatitis B (Hep B); one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (Hep A); two or three rotavirus (RV); and two

CMS ID & CQM NO.	CQM TITLE	DESCRIPTION
NEW: CMS154v1 NQF 0069	Appropriate Treatment for Children with Upper Respiratory Infection (URI)	<p>influenza (flu) vaccines by their second birthday.</p> <p>Percentage of children 3 months-18 years of age who were diagnosed with upper respiratory infection (URI) and were not dispensed an antibiotic prescription on or three days after the episode.</p>
NEW: CMS136v1 NQF 0108	ADHD: Follow-Up Care for Children Prescribed Attention Deficit/Hyperactivity Disorder (ADHD) Medication	<p>Percentage of children 6-12 years of age and newly dispensed a medication for attention deficit/hyperactivity disorder (ADHD) who had appropriate follow-up care. Two rates are reported.</p> <p>a. Percentage of children who had one followup visit with a practitioner with prescribing authority during the 30-Day Initiation Phase.</p> <p>b. Percentage of children who remained on ADHD medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two additional follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.</p>
NEW: CMS2v1 NQF 0418	Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for clinical depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen.
NEW: CMS75v1	Children who have dental decay or cavities	Percentage of children ages 0-20, who have had tooth decay or cavities during the measurement period.